

AD-A137 565

STATIONARY STOCHASTIC SSYMMETRIC CONTROL(U) STANFORD
UNIV CA DEPT OF STATISTICS H WEINER 18 JAN 84 TR-341
N00014-76-C-0475


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

(12)

ADA 137565

STATIONARY STOCHASTIC ASYMMETRIC
CONTROL

BY

HOWARD WEINER

TECHNICAL REPORT NO. 341
JANUARY 18, 1984

PREPARED UNDER CONTRACT
N00014-76-C-0475 (NR-042-267)
FOR THE OFFICE OF NAVAL RESEARCH

Reproduction in Whole or in Part is
permitted for the purpose of the United States

for public release; distribution

DEPARTMENT OF THE ARMY
TRANSFORMER GROUP
STANFORD, CALIF.

DTIC

FEB 06 1984

DTIC FILE COPY

84 02 06 014

STATIONARY STOCHASTIC ASYMMETRIC
CONTROL

BY

HOWARD WEINER

TECHNICAL REPORT NO. 341
JANUARY 18, 1984

Prepared Under Contract
N00014-76-C-0475 (NR-042-267)
For the Office of Naval Research

Herbert Solomon, Project Director


Reproduction in Whole or in Part is Permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Stationary Stochastic Asymmetric Control

by

Howard Weiner

University of California, Davis

and

Stanford University

1. Introduction Let the process $X(t)$, be defined by

$$dX(t) = u(X(t))dt + dW(t)$$

where $W(t)$ is a standard Wiener process and u is a control function, and $X(0) = x$.

Let φ be even, convex, symmetric positive, exponentially bounded and strictly increasing on the positive axis. Let the average expected cost function be

$$J(x, u) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T E [\varphi(X(s)) + |u(X(s))|^\alpha] ds$$

with $\alpha > 0$.

The object is to find the control law u which minimizes J subject to $a < U < b$ where $a < 0 < b$ are real numbers. A two-dimensional version is indicated. The case $\alpha = 1$ is considered in great detail in [1]. The results for the cases $\alpha \neq 1$ are given in this paper. The complete proofs are as in [1] and reference is made to that paper for the complete proofs of the results mentioned here. In fact, it will

suffice to indicate the nature of the solutions to the "asymptotic dynamic programming equation" in one-or multi-dimensions in [1], since the other arguments are as given there, or sufficiently similar so as to not be explicitly given. In one dimension it has the form

$$\lambda = \frac{f''}{2} + h(f') + \varphi,$$

and yields the unique optimal control by the methods of [1].

II. Optimal Law

Lemma 1. Let $\alpha = 2n$, for $n \geq 1$ an integer. Let $a < 0 < b$.

$$\text{Let } h(c) = \min_{a < u < b} (uc + u^{2n})$$

Then

$$h(c) = \begin{cases} c \left(\frac{-c}{2n} \right)^{\frac{1}{2n-1}} \left(\frac{2n-1}{2n} \right) & \text{if } \left(\frac{-c}{2n} \right)^{\frac{1}{2n-1}} \leq b \\ 0 & \text{if } c = 0 \\ \min(ac + a^{2n}, bc + b^{2n}) & \text{otherwise} \end{cases} \quad (3)$$

Proof: The proof follows by an elementary computation, noting that the minimum is either at an interior (differentiable) point or at a boundary of u .

Lemma 2. If $\alpha \neq 2n$, for some integer $n \geq 1$, for $a < 0 < b$, if

$$h(c) = \min_{a < u < b} (uc + |u|^\alpha),$$

then

$$h(c) = \begin{cases} c \left(\frac{\alpha-1}{\alpha} \right) \left(\frac{|c|}{\alpha} \right)^{\frac{1}{\alpha-1}} & \text{if } c < 0 \text{ and } \left(\frac{|c|}{\alpha} \right)^{\frac{1}{\alpha-1}} \leq b \\ \min \left(0, ac + |a|^\alpha, bc + |b|^\alpha \right) & \text{otherwise} \end{cases} \quad (4)$$

Proof The minimum is either at an interior differentiable point or at one of the three points 0, a, b.

Theorem 1 Assume $a < 0 < b$.

(a) If $d = 2n, n \geq 1$

Then there are distinct numbers $x_1 < x_2$ such that for $x_1 \leq x \leq x_2$, the "asymptotic dynamic programming equation [1], is

$$\lambda = \frac{1}{2} f'' + \left[\frac{(-f)^{2n}}{2n} \right]^{\frac{1}{2n-1}} \left(\frac{2n+1}{2n} \right) + \varphi$$

For $x > x_2$

$$\lambda = \frac{f''}{2} + af' + a^{2n} + \varphi. \quad (5)$$

The x_1, x_2, λ are found by continuity at the boundaries and by setting $f'(x_2) = f'(\infty)$ or $f'(x_1) = f'(-\infty)$.

(b) If $\alpha \neq 2n, \alpha \neq 1$

Then there are distinct numbers y_l , $1 \leq l \leq 3$

$y_1 < 0 < y_2 < y_3$, such that the "asymptotic dynamic programming equations [1]" satisfy, if

$$x \leq y_1,$$

$$\lambda = \frac{f''}{2} + bf' + |b|^\alpha + \varphi$$

and if

$$y_1 < x < 0,$$

then

$$\lambda = \frac{f''}{2} + af' + |a|^\alpha + \varphi$$

and if

$$0 < x < y_2$$

then

$$\lambda = \frac{f''}{2} + \left(\frac{(-f')^\alpha}{\alpha} \right)^{\frac{1}{\alpha-1}} \left(\frac{\alpha+1}{\alpha} \right) + \varphi$$

and if

$$y_2 < x$$

then

$$\lambda = \frac{f''}{2} + af' + |a|^\alpha + \varphi, \quad (6)$$

where the values y_1, y_2, y_3, λ are obtained by setting the solutions equal (by continuity) at y_1, y_2, y_3 and $f(y_3) = f(\infty)$ or $f(y_1) = f(-\infty)$.

Proof The expressions (5), (6) follow from (1), (2) using monotonicity and symmetry of φ , the properties of $c u + |u|^\alpha$ as a function of c, u , the consequent symmetry of f' , and the uniqueness arguments in [1].

Theorem 2

(a) If $\alpha = 2n$, if $a < 0 < b$,

the optimal law is

$$u(X) = \begin{cases} \left(-\frac{f'(X)}{2n}\right)^{\frac{1}{2n-1}} & , \quad x_1 < X < x_2 \\ b & , \quad X < x_1 \\ a & , \quad X > x_2 \end{cases} \quad (7)$$

(b) If $0 < \alpha \neq 1$, $\alpha \neq 2n$, then for $a < 0 < b$,

$$u(X) = \begin{cases} \left(-\frac{f'(X)}{\alpha}\right)^{\frac{1}{\alpha-1}} & \text{if } 0 < X < y_2 \\ b & \text{if } X < y_1 \\ a & \text{if } y_1 < X < 0 \text{ or } y_2 < X \end{cases} \quad (8)$$

Proof This follows immediately from Theorem 2 and the lemmas, (3) - (6).

Details are as in [1] for uniqueness and optimality of u .

3. Multi-dimensional case-Remarks

Let $\underline{X}(t)$ be a $k \times 1$ stochastic process with

$$d\underline{X}(t) = \underline{u}(\underline{X}(t))dt + d\underline{W}(t) \quad (9)$$

where \underline{u} is $d \times 1$ and \underline{W} is $d \times 1$ Wiener process with independent components and $\underline{X}(0) = \underline{x}$.

Let

$$J(\underline{x}, \underline{u}) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \mathbb{E} \left[\varphi(\underline{X}(s)) + \|\underline{u}(\underline{X}(s))\|_p \right] ds \quad (10)$$

where φ is even, convex, positive and exponentially bounded in each argument, and

where $\|u\|_p = \left(\sum_{\ell=1}^d |u_\ell|^p \right)^{\frac{1}{p}}$ for some $p > 0$.

It is desired to minimize J subject to

$$a_\ell \leq u_\ell \leq b_\ell \quad 1 \leq \ell \leq d$$

where $a_\ell, b_\ell, 1 \leq \ell \leq d$ are real numbers. This constraint will be written $\underline{a} \leq \underline{u} \leq \underline{b}$.

Let $\underline{c} = (c_1, \dots, c_d)$ and denote

$$h(\underline{c}) \equiv \min_{\underline{a} \leq \underline{u} \leq \underline{b}} (\underline{c} \cdot \underline{u} + \|\underline{u}\|_p) . \quad (11)$$

The evaluation of $h(\underline{c})$ requires the evaluation of $\underline{c} \cdot \underline{u} + \|\underline{u}\|_p$ at the $2^d + 1$ points given by $u_\ell = a_\ell$ or $b_\ell, 1 \leq \ell \leq d$ and at the \underline{u} minimizing $\underline{c} \cdot \underline{u} + \|\underline{u}\|_p$, for fixed \underline{c} . As \underline{c} varies, R^d is cut into regions defined by the minimum $h(\underline{c})$ and the \underline{u} achieving that minimum.

Let $\underline{f} = (f^{(1)}(\underline{x}), \dots, f^{(d)}(\underline{x}))$ for $\underline{x} = (x_1, \dots, x_d)$

such that

$$f_{ij}(\underline{x}) = \frac{\partial^2 f(\underline{x})}{\partial x_i \partial x_j} \text{ exists and is continuous } 1 \leq i, j \leq d .$$

$$\text{Let } f_j(\underline{x}) = \frac{\partial f(\underline{x})}{\partial x_j} .$$

For the various regions above, it is required to solve (for minimum λ)

$$\lambda = \frac{1}{2} \sum f_{ij}(\underline{x}) + h(f_1(\underline{x}), \dots, f_d(\underline{x})) + \varphi(\underline{x}) . \quad (12)$$

Equating solutions at the boundary of these regions defines a finite number of vectors $\underline{x}_1 = (x_{11}, \dots, x_{1d})$, $\underline{x}_2, \dots, \underline{x}_n$ where $n = 2^d + 1$ such that $f_j(\underline{x}_r) = f_j(\infty_r)$ where ∞_r denotes any ray in a boundary region such that at least one coordinate may be set equal to $\pm \infty$ and the point so obtained remain in the region. This determines λ also.

It follows as before that if \underline{u}_0 = optimal \underline{u} , then

$$\underline{u}_0(\underline{X}) = \text{that } \underline{u} \text{ giving } h(f_1, \dots, f_d)$$

in any of the regions given by the vectors as edges.

Remark 2 If in addition $\|\underline{u}\|_p \leq M < \infty$, and if \underline{u}_{0p} = optimal \underline{u} with this additional constraint, then with

$$\underline{u}_{0p} = \begin{cases} \underline{u}_0 & \text{if } \|\underline{u}_0\|_p \leq M \\ \frac{M}{R} \underline{u}_0 & \text{if } \|\underline{u}_0\|_p = R > M \end{cases} \quad (13)$$

which follows by a variational inequality. See [2] for a similar argument.

REFERENCES

1. Beneš, V, and Karatzas, I. (1980) Optimal stationary linear control of the Wiener process (unpublished manuscript).
2. Weiner, H. (1980). Constrained stochastic regulator control. (submitted).

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 341	2. GOVT ACCESSION NO. AD-A137565	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Stationary Stochastic Asymmetric Control		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL REPORT
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Howard Weiner		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0475
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-042-267
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics & Probability Program Code 411SP		12. REPORT DATE January 18, 1984
		13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Stationary, Wiener Process, Asymmetric, Bang-Bang, Optimal Control → This document describes a which deals with a cost function.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The one-dimensional problem considered is the minimization of the functional. $J(x,u) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T E (\varphi(X(s)) + u(X(s)) ^\alpha) ds \quad \text{where } \alpha > 0, \alpha \neq 1, \quad \text{with}$ $dX(t) = u(X(t))dt + dW(t), \quad X(0) = x, \quad \text{and } -\infty < a < U < b < \infty.$ Generalizations to a multi-dimensional case are mentioned.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

END

FILMED

3-84

DTIC